

SOLITON SOLUTIONS OF THE σ MODEL AND THE DISORIENTED CHIRAL CONDENSATES

Prasanta. K. Panigrahi¹ and C. Nagaraja Kumar

School of Physics, University of Hyderabad, Hyderabad - 500 046 (INDIA).

ABSTRACT

We derive travelling soliton solutions of the σ model, relevant for the description of dynamics of pions, in the high-energy, heavy-ion collisions. These extended objects are analogs of Neél and Bloch type defects in ferromagnetic systems and could possibly describe the disoriented chiral condensates. It is shown that these solutions are metastable and can naturally produce an asymmetry between π^0 and π^+ , π^- mesons in the soft-pion emissions from heavy-ion collisions.

The formation of the disoriented chiral condensates (DCC) in the high-energy, heavy-ion collisions and their detection through the asymmetry in the soft π^0 and π^+ , π^- emissions have generated considerable interest in the recent literature [1,2,3]. There are also experimental indications pointing to the possible formation of DCC in cosmic ray collisions[4]. Broadly speaking, DCC refers to the domains in space-time, where the chiral order parameter points along a direction in the iso-spin space, different from the iso-spin singlet σ -direction in the outside physical vacuum. It is by now well accepted that the σ -model with four fields σ, π^+, π^- and π^0 having an approximate $SU(2) \otimes SU(2)$ symmetry effectively captures the low-energy dynamics of QCD with two flavors. Below a certain critical temperature T_c , the global $SU(2)$ chiral symmetry is spontaneously broken due to the non-vanishing vacuum expectation value (VEV) of the iso-singlet fermion bilinear $\langle \bar{\psi}\psi \rangle$. In the σ model description, this corresponds to the σ field having a VEV and further more the pions are realized as the Goldstone modes in the broken symmetry phase. The chiral symmetry of the QCD Lagrangian is only approximate because of the presence of the current quark masses and hence the pions acquire a small but non-vanishing mass.

¹ email: panisp@uohyd.ernet.in

In high energy, heavy-ion collisions, there exists a possibility that chiral symmetry gets restored in a small region of hot hadronic matter in the central zone. This region surrounded by an outwardly expanding hot hadronic shell cools rapidly to a temperature below T_c ; the chiral order parameter in this domain can in principle point in an isospin direction different from the σ field. The eventual realignment with the physical vacuum at a later period in time can result in the emission of soft, coherent pions. There could be anomalous fluctuations in neutral to charged pion ratio. If the DCC domain is not baryon rich the dynamics can be described by the σ model. Both the linear and the non-linear σ models have been extensively used to study the formation of DCC near T_c .

In the following, we follow the non-equilibrium phase transition picture of DCC formation originally due to Rajagopal and Wilczek [5]. Here quenching results in the removal of σ and $\vec{\pi}$ fields from contact with the heat bath. These fields then evolve according to the $T = 0$ Hamiltonian. It has been shown numerically, in the context of the linear σ model that in intermediate times, $5 \sim 50 \times \frac{1}{m_\pi}$, there is a dramatic amplification of long wavelength modes. Since short wavelength modes are not completely absent, the resulting domains are not smooth [6].

Hence the emerging physical picture of DCC is more closer to a ‘pion laser’ where the order parameter instead of being a constant throughout the DCC can fluctuate and hence is space-time dependent. The nonlinearity of the σ model makes it difficult to get the analytic time evolution in $(3 + 1)$ dimensions. Furthermore mean field approximation may not be fully reliable for the entire intermediate time range. Keeping these in mind, we start with the idealized Heisenberg type boundary condition : the thin disc representing the Lorentz-contracted nuclei at the time of collision is infinite in extent in the transverse direction . Assuming the fields to be independent of the transverse directions one can simplify the problem to a $(1 + 1)$ dimensional field theory [7] . Inspired by effective field theories describing lasers [8] and other non-equilibrium phenomena [9] we consider the σ model Lagrangian in the presence of additional isospin violating quadratic terms in the field variables and work out the consequences. These terms could possibly arise due to non-equilibrium effects or squeezing [10]. The relevant

Lagrangian then is

$$\mathcal{L} = \frac{f_\pi^2}{2}[(\partial_\mu \sigma)^2 + (\partial_\mu \pi^a)^2] - V(\pi^a, \sigma) \quad (1)$$

$$V(\pi^a, \sigma) = \lambda\{\sigma^2 + \pi^{a2} - 1\}^2 - \frac{\mu}{2}\pi_3^2 + \kappa \quad (2)$$

κ was added to adjust the lowest value of the potential and the term with coefficient μ is the new term. The $\pi_\pm$ are $\frac{1}{\sqrt{2}}(\pi_1 \pm i\pi_2)$. It is easy to solve for the solutions for the equations of motion and we find

$$\pi_1 = l \operatorname{sech} \frac{\sqrt{2\mu}}{f_\pi} \xi \quad (3)$$

$$\pi_2 = m \operatorname{sech} \frac{\sqrt{2\mu}}{f_\pi} \xi \quad (4)$$

$$\pi_3 = \alpha \tanh \frac{\sqrt{2\mu}}{f_\pi} \xi \quad (5)$$

$$\sigma = n \operatorname{sech} \frac{\sqrt{2\mu}}{f_\pi} \xi \quad (6)$$

$$\text{where} \quad \alpha^2 = 1 + \frac{\mu}{2\lambda} \quad (7)$$

$$\text{and} \quad l^2 + m^2 + n^2 = 1 - \frac{\mu}{2\lambda} \quad (8)$$

this last relation puts a restriction on the values μ can take to be less than 2λ . The integrated energy density for large distance L is given by

$$\sqrt{\frac{\mu}{2}} f_\pi \left[2 - \frac{\mu}{3\lambda} \right] \quad (9)$$

The ratio of the number of neutral pions to the total pions produced is proportional to ratio of the squared amplitudes of the corresponding pion fields integrated over the length L .

$$\frac{N(\pi_0)}{N_{\text{total}}} = \frac{\int \pi_3^2 dx}{\int (\pi_a^2 + \sigma^2) dx} = \frac{\alpha^2 (L - \frac{\operatorname{Tanh}(aL)}{a})}{(\alpha^2 L - \frac{\mu}{\lambda} \frac{\operatorname{Tanh}(aL)}{a})} \quad (10)$$

where a is $\frac{\sqrt{2\mu}}{f_\pi}$. For small L this ratio tends to zero and for large distances it gives the value 1.

Another interesting term which can be added is $\mu\pi_1^2 = \mu\frac{1}{2}(\pi_+ + \pi_-)^2$. This term leads to Bloch type solitons given by $A \operatorname{Tanh}(m\xi) + iB \operatorname{Sech}(m\xi)$ which smoothly connects π_+ with π_- asymptotically. However, this term

violates $U(1)$ electromagnetic invariance and hence may not be relevant in the present situation.

In conclusion, the soliton solutions which appear in other fields of non-equilibrium phenomena can possibly play a significant role in the DCC formation. A number of investigations e.g., stability analysis, coupling with fermions needs to be carried out for illustrating their role [11]. A first principle study of the origin of the additional terms starting from non-equilibrium field theory also needs to be carried out.

Acknowledgments: One of us (P.K.P) would like to thank Drs. R. MacKenzie and M. Paranjape for hospitality and Drs. K. Rajagopal and T.Cohen for many discussions and clarifications. (C.N.K) research is supported by CSIR, India, through S.R.A. Scheme.

References

- 1 F. Wilczek, Int. Journ. Mod. Phys. **A7** , 3911 (1992), J.Bjorken, Int. J. Mod. Phys. **A7**, 4189 (1992)
- 2 K.Rajagopal and F. Wilczek, Nucl. Phys. **B399**, 395 (1993).
- 3 A.Anslem, Phys. Lett. **B 217** 169, (1988).
- 4 Chacaltaya-Pamir Collaboration; Tokyo University preprint ICRR - Report -258-91-27 and references there in
- 5 K. Rajagopal, preprint, hep-ph 9703258, J.D. Bjorken, K.L.Kowalski and C.C.Taylor SLAC Preprint SLAC -Pub-6109.
- 6 G. Amelino-Camelia, J.D. Bjorken and S.E. Larsson, preprint SLAC-PUB-7565, hep-ph 9706530.
- 7 J.-P. Blaizot and A. Krzywicki, Phys. Rev. **D 46** 246, (1992)
- 8 S.Longhi, Europhysics Letters **37**, 257, (1997).
- 9 P. Coulet and K. Emilsson Physica **D 61** 119, (1992).
- 10 B.A. Bambah, preprint, hep-ph 9708414 and references therein.
- 11 P.K.Panigrahi and C.N.Kumar, manuscript under preparation.